

Orbital effect of Cooper pairs on Kondo effect in unconventional superconductors

Masashige Matsumoto¹ and Mikito Koga²

¹*Department of Physics, Faculty of Science, Shizuoka University, 836 Oya, Shizuoka 422-8529, Japan*

²*Department of Physics, Faculty of Education, Shizuoka University, 836 Oya, Shizuoka 422-8529, Japan*

(March 3, 2001)

A new type of Kondo effect peculiar to unconventional superconductors is studied theoretically by using the Wilson's numerical renormalization group method. In this case, an angular momentum of a Cooper pair plays an important role in the Kondo effect. It produces multichannel exchange couplings with a local spin. Here we focus on a $p_x + ip_y$ -wave state which is a full gap system. The calculated impurity susceptibility shows that the local spin is almost quenched by the Kondo effect in the strong coupling region ($T_K/\Delta \rightarrow \infty$), while the ground state is always a spin doublet over all the T_K/Δ region. Here T_K and Δ are the Kondo temperature and the superconducting energy gap, respectively. This is different from the s -wave pairing case where the Kondo singlet is realized for large T_K/Δ values. The strong coupling analysis shows that the $p_x + ip_y$ -wave Cooper pair is connected to the Kondo singlet via the orbital dynamics of the paired electrons, generating the spin of the ground state. This type of Kondo effect reflects the symmetry of the conduction electron system.

I. INTRODUCTION

The search for variety in the Kondo effect was stimulated by the discovery of the non-Fermi liquid behavior in heavy fermion materials and by the observation of the two-level system character in metallic glasses and nanoscale devices. In these cases, orbital degrees of freedom are very important as well as spin.^{1,2} Due to incomplete compensation of a local moment, it displays singular behavior in physical quantities, differing from the ordinary single-channel Kondo effect. This Kondo problem has been investigated mainly for metallic systems with no gap in the density of states.

For a gapped system, the problem of a magnetic impurity in a s -wave superconductor has been investigated for a long time.³⁻¹⁰ Since the s -wave property is described by a single channel, the Kondo effect itself is understood in the framework of the conventional theory based on the single-channel Kondo model. On the other hand, no one has paid attention to multichannel properties of p -wave and d -wave superconductors derived from angular momenta of their Cooper pairs. They can exhibit a new type of Kondo effect peculiar to the unconventional superconductivity.

The unconventional superconductivity is very sensitive to impurities and surface boundaries. It displays peculiar phenomena not observed in the s -wave superconductivity. It is known that scattering objects destroy the unconventional superconductivity and induce bound states.¹¹ The nuclear magnetic resonance (NMR) T_1^{-1} measurement performed in Zn-doped $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ showed a constant T_1T at low temperatures¹² reflecting finite density of states at lower energies. As shown by the low temperature scanning tunnel microscope (STM), a single Zn atom actually generates the low energy bound state.¹³ For the $d_{x^2-y^2}$ -wave, the (1,1,0) surface is regarded as the pair-breaking boundary.¹⁴ A very recent experiment showed a zero energy peak in the tunneling conductance

between the (1,1,0) boundary of $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ and a normal metal.¹⁵ Thus much attention has been paid to the effects of non-magnetic impurities and static boundaries in the last decade.

Recently the effect of magnetic impurities in unconventional superconductors has attracted attention gradually with the development of experimental studies. According to the NMR experiments,¹⁶ a local moment is induced around a Zn site substituted for a Cu site in $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$. It is expected that the observed Kondo behavior is due to this moment.¹⁷ Similar behavior is also observed in Li-substituted $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$.¹⁸ In this case, the local magnetic susceptibility of the Li induced moment shows strong reduction of the Kondo screening below the superconducting transition temperature. At present the search for the influence of unconventional superconductivity on the Kondo effect is still in progress. We are also interested in the Kondo effect for different types of superconductors other than such high- T_c cuprates.

Several studies of a magnetic impurity in a conventional (s -wave) superconductor clarified how a bound state generated around the impurity depends on the Kondo temperature T_K and the superconducting energy gap Δ .⁶⁻¹⁰ The numerical renormalization group (NRG) study by Satori *et al.*⁸ showed that the bound state energy is scaled by T_K/Δ and the interchange of singlet and doublet ground states occurs smoothly. Even if an energy gap exists in the density of quasiparticle states, a local spin couples with the quasiparticles antiferromagnetically to form a spin singlet (Kondo singlet) for $T_K/\Delta > 0.3$. A similar competition was discussed in gapless $d_{x^2-y^2}$ -wave superconductors such as high T_c cuprates.¹⁹ In this case, the gapless density of states stabilizes the Kondo singlet more favorably than in the gapped s -wave case. However, it is not certain that a singlet ground state is always realized in all types of superconductors when the Kondo effect is strong enough. A Cooper pair in the unconventional superconductivity has an angular momentum and

can produce multichannel scattering of quasiparticles at an impurity. As well as for metals, a difference between single channel and multichannel properties in the Kondo effect can be expected for superconductors.

In order to elucidate the multichannel property, we study the Kondo effect due to a local spin coupled to a $p_x + ip_y$ (or $d_{x^2-y^2} + id_{xy}$)-wave superconducting state. The $\mathbf{d}$ -vector for the $p_x + ip_y$ -wave is given by $\mathbf{d}_{\mathbf{k}} = \hat{z}(k_x + ik_y)$ where $\hat{z}$ is a unit vector in the z axis. Since the density of states has a full gap like the s -wave case, we can find the peculiarity of the unconventional superconductivity in comparison with the s -wave. The property of the $p_x + ip_y$ superconductivity is analogous to that of the A phase found in the superfluid ^3He . A lot of attention has been paid to this superconductivity since the discovery of the spin triplet superconductor Sr_2RuO_4 .^{20–22} The recent theoretical studies proposed a number of unique physics associated with the broken time reversal symmetry observed by the μSR experiment.²³ In the $p_x + ip_y$ -wave superconductor, spontaneous current appears close to the scattering objects and it induces a finite magnetic field.^{24,25} This argument explains the μSR data well.²⁶ If the system induces the spontaneous magnetic field by itself, a spontaneous Hall effect can be expected.²⁷ A vortex charging effect is also interesting, peculiar to the rotating Cooper pairs.²⁸

The most important point in this paper is that for the $p_x + ip_y$ -wave case one of the Cooper pairing electrons is coupled to a local spin and the other has no direct coupling with the local spin. As we will discuss later, the latter can interact with the local spin through the superconducting order parameter. We use the NRG method^{29–31} to study this new type of two-channel Kondo effect. As for the s -wave case, we discretize the bare conduction band logarithmically and apply the Bogoliubov transformation to obtain a diagonal form of NRG Hamiltonian for the superconductivity. In the NRG Hamiltonian, the two independent channels consisting of the $l = 0$ and $l = 1$ orbitals are coupled to each other at the impurity. The NRG result shows that the ground state is a spin doublet over all the T_K/Δ region and T_K/Δ dependence of a bound state expresses a competition between the Kondo effect and the superconducting energy gap. The solution of the strong coupling limit ($T_K/\Delta \rightarrow \infty$) shows that the ground state is a spin doublet due to dynamical coupling between the $l = 0$ and $l = 1$ electrons, while the local spin is almost quenched by the $l = 0$ electrons.

This paper is organized as follows. In Sec. II we derive the NRG Hamiltonian to investigate the two-channel Kondo effect specific to the $p_x + ip_y$ (or $d_{x^2-y^2} + id_{xy}$)-wave superconductivity. Here the origin of the two channels is explained thoroughly. In Sec. III we show the NRG results in comparison with the s -wave case. We also calculate a local spin susceptibility and discuss the Ising case. It is very useful to understand the quantum effect of the magnetic impurity in the gapped electron system. This paper is closed with concluding remarks in Sec. IV.

II. FORMULATION

In this section we derive the NRG Hamiltonian for the Kondo effect in the $p_x + ip_y$ -wave ($d_{x^2-y^2} + id_{xy}$ -wave) superconductor. Here we study a single magnetic impurity at the center in two-dimensional superconducting systems.

A. Model Hamiltonian

Let us begin with the following Hamiltonian:

$$H = H_{\text{kin}} + H_{\Delta} + H_{\text{imp}}, \quad (2.1a)$$

$$H_{\text{kin}} = \sum_{\mathbf{k}\sigma} \epsilon_{\mathbf{k}} a_{\mathbf{k}\sigma}^{\dagger} a_{\mathbf{k}\sigma}, \quad (2.1b)$$

$$H_{\Delta} = \sum_{\mathbf{k}} \left(\Delta_{\mathbf{k}} a_{\mathbf{k}\uparrow}^{\dagger} a_{-\mathbf{k}\downarrow}^{\dagger} + \text{H.c.} \right), \quad (2.1c)$$

$$H_{\text{imp}} = \frac{1}{2} \sum_{\mathbf{k}\mathbf{k}'\sigma\sigma'} (-J\mathbf{S} \cdot \boldsymbol{\sigma}_{\sigma\sigma'} + V\delta_{\sigma,\sigma'}) a_{\mathbf{k}\sigma}^{\dagger} a_{\mathbf{k}'\sigma'}. \quad (2.1d)$$

Here $\mathbf{k}$ and σ represent the wave vector and spin of the conduction electrons, respectively. $a_{\mathbf{k}\sigma}$ is the annihilation operator of the conduction electron, $\epsilon_{\mathbf{k}}$ is the kinetic energy of the conduction electron, and $\Delta_{\mathbf{k}}$ is the momentum dependent order parameter. $\mathbf{S}$ expresses the $S = 1/2$ spin operator for the magnetic impurity, and $\boldsymbol{\sigma}$ is the Pauli matrix for the conduction electrons. $J(< 0)$ and V are the antiferromagnetic and non-magnetic couplings, respectively. Here we have assumed a short-range (s -wave) scattering impurity. For the $p_x + ip_y$ -wave pairing state, the order parameter has a form of $\Delta_{\mathbf{k}} = \Delta e^{i\phi_{\mathbf{k}}}$. Here Δ is the order parameter which we assume a real number, and $\phi_{\mathbf{k}}$ is the angle of the wave vector measured from the k_x axis. Since the orbital angular momentum is a good quantum number for the $p_x + ip_y$ -wave, it is convenient to transform the wave vector representation into the following two-dimensional polar coordinate representation:

$$a_{\mathbf{k}\sigma} = \sum_l (-i)^l |J_{l+1}(kR)| e^{il\phi_{\mathbf{k}}} a_{kl\sigma}, \quad (2.2)$$

where J_l is the l -th Bessel function, and R is the radius of the two-dimensional system. l is the z -component of the orbital angular momentum of the conduction electron. Substituting Eq. (2.2) into Eq. (2.1), we obtain

$$H = B (\bar{H}_{\text{kin}} + \bar{H}_{\Delta} + \bar{H}_{\text{imp}}), \quad (2.3a)$$

$$\bar{H}_{\text{kin}} = \int_{-1}^1 dk k \sum_{l\sigma} a_{kl\sigma}^{\dagger} a_{kl\sigma}, \quad (2.3b)$$

$$\bar{H}_{\Delta} = \int_{-1}^1 dk \sum_l (-1)^{1-l} \left(i\bar{\Delta} a_{kl\uparrow}^{\dagger} a_{k,-l+1,\downarrow}^{\dagger} + \text{H.c.} \right), \quad (2.3c)$$

$$\bar{H}_{\text{imp}} = \sum_{\sigma\sigma'} (-\bar{J}\mathbf{S} \cdot \boldsymbol{\sigma}_{\sigma\sigma'} + \bar{V}\delta_{\sigma,\sigma'}) f_{00\sigma}^{\dagger} f_{00\sigma'}, \quad (2.3d)$$

$$f_{00\sigma} = \frac{1}{\sqrt{2}} \int_{-1}^1 dk a_{k,l=0,\sigma}, \quad (2.3e)$$

where $B = k_F^3 R / (\pi m) = 2k_F R E_F / \pi$, $\bar{\Delta} = \Delta / (2E_F)$, $\bar{J} = J\rho$, $\bar{V} = V\rho$. Here $E_F = k_F^2 / (2m)$ is the Fermi energy, and $\rho = mR^2 / 2 = (k_F R)^2 / (4E_F)$ is the density of states at the Fermi energy for the two-dimensional conduction band. In Eq. (2.3) we have used the following asymptotic form of the Bessel function: $J_l(kR) \simeq \sqrt{2/(\pi kR)}$ (for a large kR). The wave number k in Eq. (2.3) is normalized by k_F . The kinetic energy of the conduction electron is linearized near the Fermi energy. For simplicity, we have used the same values for both the superconducting cutoff energy and the band width k_F , since it is assured that this does not alter the results.⁹ The fermion represented by the operator $f_{00\sigma}$ corresponds to a fermion localized at the impurity. The first subscript 0 of $f_{00\sigma}$ represents the 0-th shell of the NRG Hamiltonian which will be derived later (see Appendix). The second zero means the $l = 0$ orbital. Notice that the angular momentum of the $p_x + ip_y$ -wave Cooper pair equals to one (see $\bar{H}_\Delta$), and that the $l = 0$ and $l = 1$ orbitals are coupled. Since the angular momentum is a good quantum number for the $p_x + ip_y$ -wave, both orbitals are decoupled from the other orbitals. Due to the short-range (s -wave) scattering, the electrons only in the $l = 0$ orbital couple with the impurity. The electrons in the $l = 1$ orbital can also interact with the impurity via the $p_x + ip_y$ -wave order parameter. It is sufficient to choose only the $l = 0$ and $l = 1$ orbitals in Eq. (2.3) to study the Kondo effect for the $p_x + ip_y$ -wave state. Therefore we can apply the NRG method to this Kondo problem as we do to the two-channel Kondo problem.^{32,33} We rewrite then $\bar{H}_{\text{kin}}$ and $\bar{H}_\Delta$ as

$$\bar{H}_{\text{kin}} = \int_{-1}^1 dk k \sum_{\sigma} \left(a_{k0\sigma}^\dagger a_{k0\sigma} + a_{k1\sigma}^\dagger a_{k1\sigma} \right), \quad (2.4a)$$

$$\bar{H}_\Delta = \int_{-1}^1 dk \sum_{\sigma} \left(i\bar{\Delta} a_{k1\sigma}^\dagger a_{k0,-\sigma}^\dagger + \text{H.c.} \right). \quad (2.4b)$$

We can obtain a similar $\bar{H}_\Delta$ for a $d_{x^2-y^2} + id_{xy}$ -wave if we replace $i\bar{\Delta}$ by $-\sigma\bar{\Delta}$ and $l = 1$ by $l = 2$, since the order parameter for the $d_{x^2-y^2} + id_{xy}$ -wave has a form of $\Delta_{\mathbf{k}} = \Delta e^{i2\phi_{\mathbf{k}}}$.

B. Transformation of the Hamiltonian

In our formulation we follow the procedure by Sakai *et al.*⁹ and derive a NRG Hamiltonian. There is another way to derive the Hamiltonian,⁸ which we describe in Appendix. First we diagonalize the conduction electron part by introducing the following Bogoliubov transformation:

$$a_{k0\sigma} = u_k \alpha_{k,+,\sigma} - v_k^* \alpha_{k,-,-\sigma}^\dagger, \quad (2.5a)$$

$$a_{k1\sigma} = u_k \alpha_{k,-,\sigma} + v_k^* \alpha_{k,+,-\sigma}^\dagger. \quad (2.5b)$$

Here $\alpha_{k,\tau=\pm,\sigma}$ is the annihilation operator for the superconducting quasiparticle with σ spin in τ channel. The coefficients u_k and v_k are defined by

$$u_k = \sqrt{\frac{1}{2} \left(1 + \frac{|k|}{E_k} \right)}, \quad (2.6a)$$

$$v_k = i \sqrt{\frac{1}{2} \left(1 - \frac{|k|}{E_k} \right)} \text{sgn}(k), \quad (2.6b)$$

where $E_k = (k^2 + \bar{\Delta}^2)^{1/2}$ is the energy of the quasiparticles. The $p_x + ip_y$ -wave Hamiltonian is then expressed in terms of the superconducting quasiparticle as

$$\begin{aligned} \bar{H}_{p_x+ip_y} &= \bar{H}_{\text{kin}} + \bar{H}_\Delta \\ &= \int_{-1}^1 dk E_k \text{sgn}(k) \sum_{\tau=\pm,\sigma} \alpha_{k\tau\sigma}^\dagger \alpha_{k\tau\sigma}. \end{aligned} \quad (2.7)$$

For the later convenience, we introduce the following transformations:

$$\beta_{|k|,-\tau,-\sigma}^\dagger = \alpha_{-|k|,\tau,\sigma}, \quad (2.8a)$$

$$A_{k\tau\sigma} = \frac{1}{\sqrt{2}} (\alpha_{k\tau\sigma} - i\tau\beta_{k\tau\sigma}), \quad (2.8b)$$

$$B_{k\tau\sigma} = \frac{1}{\sqrt{2}} (\alpha_{k\tau\sigma} + i\tau\beta_{k\tau\sigma}), \quad (2.8c)$$

$$C_{+,\sigma}(k) = A_{k,+,\sigma}, \quad (2.8d)$$

$$C_{+,\sigma}(-k) = -iB_{k,-,-\sigma}^\dagger, \quad (2.8e)$$

$$C_{-,\sigma}(k) = iA_{k,-,-\sigma}^\dagger, \quad (2.8f)$$

$$C_{-,\sigma}(-k) = B_{k,+,\sigma}. \quad (2.8g)$$

The Hamiltonian (2.7) can be expressed by the above C operators as

$$\bar{H}_{p_x+ip_y} = \int_0^1 dk E_k \sum_{s=\pm,\tau,\sigma} \tau s C_{\tau\sigma}^\dagger(sk) C_{\tau\sigma}(sk). \quad (2.9)$$

For the impurity part, the operator $f_{00\sigma}$ is also expressed in terms of the superconducting quasiparticle as

$$\begin{aligned} f_{00\sigma} &= \frac{1}{\sqrt{2}} \int_0^1 dk \left[u_k (\alpha_{k+\sigma} + \beta_{k,-,-\sigma}^\dagger) \right. \\ &\quad \left. - v_k^* (\alpha_{k,-,-\sigma}^\dagger - \beta_{k+,\sigma}) \right] \\ &= \frac{1}{\sqrt{2}} \int_0^1 dk \left[w_{k+} (A_{k+\sigma} + iA_{k,-,-\sigma}^\dagger) \right. \\ &\quad \left. + w_{k-} (B_{k+\sigma} - iB_{k,-,-\sigma}^\dagger) \right], \end{aligned} \quad (2.10a)$$

$$w_{k,\pm} = \sqrt{\frac{1}{2} \left(1 \pm \frac{|\bar{\Delta}|}{E_k} \right)}. \quad (2.10b)$$

By introducing the following operators

$$a_{\tau,\sigma} = N_+^{-1} \int_0^1 dk w_{k+} A_{k,\tau,\sigma}, \quad (2.11a)$$

$$b_{\tau,\sigma} = N_-^{-1} \int_0^1 dk w_{k-} B_{k,\tau,\sigma}, \quad (2.11b)$$

$$N_{\pm} = \left(\int_0^1 dk w_{k\pm}^2 \right)^{1/2}, \quad (2.11c)$$

$f_{00\sigma}$ is expressed as

$$\begin{aligned} f_{00\sigma} &= \frac{1}{\sqrt{2}} \left[N_+ a_{+,\sigma} - i N_- b_{-,\sigma}^\dagger \right. \\ &\quad \left. + i (N_+ a_{-,\sigma}^\dagger - i N_- b_{+,\sigma}^\dagger) \right] \\ &= \frac{1}{\sqrt{2}} (c_{0,+,\sigma} + c_{0,-,\sigma}), \end{aligned} \quad (2.12)$$

where $c_{0,\tau,\sigma}$ is defined by

$$\begin{aligned} c_{0,+,\sigma} &= N_+ a_{+,\sigma} - i N_- b_{-,\sigma}^\dagger \\ &= \int_0^1 dk \left(w_{k+} A_{k,+,\sigma} - i w_{k-} B_{k,-,\sigma}^\dagger \right) \\ &= \int_0^1 dk \sum_{s=\pm} w_{k,s} C_{+,\sigma}(sk), \end{aligned} \quad (2.13a)$$

$$\begin{aligned} c_{0,-,\sigma} &= i \left(N_+ a_{-,\sigma}^\dagger - i N_- b_{+,\sigma}^\dagger \right) \\ &= \int_0^1 dk \left(w_{k+} i A_{k,-,\sigma}^\dagger + w_{k-} B_{k,+,\sigma} \right) \\ &= \int_0^1 dk \sum_{s=\pm} w_{k,s} C_{-,\sigma}(sk). \end{aligned} \quad (2.13b)$$

The exchange Hamiltonian $\bar{H}_{\text{imp}}$ is then written as

$$\bar{H}_{\text{imp}} = \frac{1}{2} \sum_{\tau\tau'\sigma\sigma'} (-\bar{J} \mathbf{S} \cdot \boldsymbol{\sigma}_{\sigma\sigma'} + \bar{V} \delta_{\sigma,\sigma'}) c_{0\tau\sigma}^\dagger c_{0\tau'\sigma'}. \quad (2.14)$$

In Eq. (2.14), the subscript $\tau = \pm$ represents the two channels, both of which consists of the $l = 0$ and $l = 1$ orbitals. While the impurity interacts with only the $l = 0$ electron, Eq. (2.14) contains the channel-flip ($\sum_{\tau\tau'}$) terms. This is because the operator $f_{00\sigma}$ is expressed as Eq. (2.12). The two-channel property in Eq. (2.14) originates from the angular momentum of the $p_x + ip_y$ -wave Cooper pair. In order to treat the Kondo problem in the superconducting state, first we have diagonalized the Hamiltonian of the conduction electron part. The diagonalized representation (τ representation) in this Hamiltonian is given by the linear combination of the $l = 0$ and $l = 1$ orbitals, since the $p_x + ip_y$ -wave mixes the two orbitals. It is natural that the impurity part of the Hamiltonian contains the channel-flip part in the τ representation.

In contrast to the present $p_x + ip_y$ -wave case, the standard two-channel Kondo model was derived from the internal degrees of freedom of the impurity atom.^{1,34} In our

model, the impurity does not have such internal degrees of freedom except for the $S = 1/2$ spin. However, we can obtain the two-channel property due to the orbital effect of the Cooper pairs. Therefore the present two-channel property for the $p_x + ip_y$ -wave can provide new Kondo physics peculiar to the unconventional superconductors.

C. Discretization

Following Wilson's procedure,²⁹ we discretize the conduction band with a logarithmic mesh in the k space. The Hamiltonian for the discretized conduction band is written as

$$\bar{H}_{p_x + ip_y} = \sum_{m=0}^{\infty} \sum_{\tau\sigma s} \tau s E_m C_{\tau\sigma}^\dagger(sm) C_{\tau\sigma}(sm), \quad (2.15a)$$

$$E_m = \sqrt{k_m^2 + \bar{\Delta}^2}, \quad (2.15b)$$

$$k_m = (1 + \Lambda^{-1}) \Lambda^{-m} / 2, \quad (2.15c)$$

where $\Lambda > 1$ is the discretization parameter. The above C operators satisfies the Fermion commutation relation:

$$\{C_{\tau\sigma}(sm), C_{\tau'\sigma'}^\dagger(s'm')\} = \delta_{\tau,\tau'} \delta_{\sigma,\sigma'} \delta_{s,s'} \delta_{m,m'}. \quad (2.16)$$

The fermion operator for the localized orbit (the 0-th shell orbit) $c_{0\tau\sigma}$ is also expressed with the C operators as

$$c_{0\tau\sigma} = (1 - \Lambda^{-1})^{-1/2} \sum_{sm} w_{ms} \Lambda^{-m/2} C_{\tau\sigma}(sm), \quad (2.17a)$$

$$w_{ms} = \sqrt{(1 + s \frac{\bar{\Delta}}{E_m})} / 2. \quad (2.17b)$$

We define $c_{n\tau\sigma}$ for the n -th shell orbit and satisfies the following commutation relation:

$$\begin{aligned} [c_{n\tau\sigma}, \bar{H}_{p_x + ip_y}] &= t_{n-1} c_{n-1,\tau\sigma} + \tau \eta_n c_{n\tau\sigma} \\ &\quad + t_n c_{n+1,\tau\sigma}. \end{aligned} \quad (2.18)$$

For $n = 0$ we have

$$\begin{aligned} [c_{0\tau\sigma}, \bar{H}_{p_x + ip_y}] &= (1 - \Lambda^{-1})^{1/2} \\ &\quad \times \sum_{sm} w_{ms} \Lambda^{-m/2} \tau s E_m C_{\tau\sigma}(sm) \\ &= \tau \eta_0 c_{0\tau\sigma} + t_0 c_{1\tau\sigma}, \end{aligned} \quad (2.19)$$

where η_0 , t_0 , and $c_{1\tau\sigma}$ are defined by

$$\eta_0 = (1 - \Lambda^{-1}) \sum_{sm} w_{ms}^2 \Lambda^{-m} s E_m, \quad (2.20a)$$

$$t_0 = \left[(1 - \Lambda^{-1}) \sum_{sm} w_{sm}^2 \Lambda^{-m} (s E_m - \eta_0)^2 \right]^{1/2}, \quad (2.20b)$$

$$c_{1\tau\sigma} = \tau \frac{(1 - \Lambda^{-1})^{1/2}}{t_0}$$

$$\times \sum_{sm} w_{sm} \Lambda^{-m/2} (sE_m - \eta_0) C_{\tau\sigma}(sm). \quad (2.20c)$$

Repeating this procedure, we obtain the total Hamiltonian which has the following hopping type with staggered potentials:⁹

$$\bar{H} = \bar{H}_{p_x + ip_y} + \bar{H}_{\text{imp}}, \quad (2.21a)$$

$$\begin{aligned} \bar{H}_{p_x + ip_y} = \sum_{n\tau\sigma} & \left[\tau \eta_n c_{n\tau\sigma}^\dagger c_{n\tau\sigma} \right. \\ & \left. + t_n \left(c_{n+1,\tau\sigma}^\dagger c_{n\tau\sigma} + c_{n,\tau\sigma}^\dagger c_{n+1,\tau\sigma} \right) \right], \end{aligned} \quad (2.21b)$$

$$\bar{H}_{\text{imp}} = \frac{1}{2} \sum_{\tau\tau'\sigma\sigma'} (-\tilde{J}\mathbf{S} \cdot \boldsymbol{\sigma}_{\sigma\sigma'} + \tilde{V}\delta_{\sigma,\sigma'}) c_{0\tau\sigma}^\dagger c_{0\tau'\sigma'}. \quad (2.21c)$$

Here η_n and t_n are given by

$$\eta_n = (-1)^n \bar{\Delta}, \quad (2.22a)$$

$$t_n = \frac{(1 + \Lambda^{-1})}{2} \varepsilon_n \Lambda^{-n/2}, \quad (2.22b)$$

$$\begin{aligned} \varepsilon_n = & \left[1 - \Lambda^{-(n+1)} \right] \left[1 - \Lambda^{-(2n+1)} \right]^{-1/2} \\ & \times \left[1 - \Lambda^{-(2n+3)} \right]^{-1/2}. \end{aligned} \quad (2.22c)$$

In the practical calculation, we use the following scaled Hamiltonian in a recursion relation given by

$$\begin{aligned} H_{N+1} = \Lambda^{1/2} H_N + \sum_{\tau\sigma} & \left[\varepsilon_N (c_{N+1,\tau\sigma}^\dagger c_{N\tau\sigma} + \text{H.c.}) \right. \\ & \left. + (-1)^N \Lambda^{N/2} \tau \tilde{\Delta} c_{N+1,\tau\sigma}^\dagger c_{N+1,\tau\sigma} \right], \end{aligned} \quad (2.23a)$$

$$\begin{aligned} H_0 = & \left[\frac{1}{2} \sum_{\tau\tau'\sigma\sigma'} (-\tilde{J}\mathbf{S} \cdot \boldsymbol{\sigma}_{\sigma\sigma'} + \tilde{V}\delta_{\sigma,\sigma'}) c_{0\tau\sigma}^\dagger c_{0\tau'\sigma'} \right. \\ & \left. - \sum_{\tau\sigma} \tau \tilde{\Delta} c_{0\tau\sigma}^\dagger c_{0\tau\sigma} \right] \Lambda^{-1/2}. \end{aligned} \quad (2.23b)$$

Here the parameters are defined by

$$\begin{aligned} \tilde{\Delta} &= \frac{\bar{\Delta}}{(1 + \Lambda^{-1})/2}, \\ \tilde{J} &= \frac{\bar{J}}{(1 + \Lambda^{-1})/2}, \\ \tilde{V} &= \frac{\bar{V}}{(1 + \Lambda^{-1})/2}. \end{aligned} \quad (2.24)$$

The Hamiltonian (2.21) is related to H_N and is given by

$$\bar{H} = \frac{1 + \Lambda^{-1}}{2} \lim_{N \rightarrow \infty} \Lambda^{-(N-1)/2} H_N. \quad (2.25)$$

In Eq. (2.23), $c_{N\tau\sigma}$ is an operator of the NRG fermion quasiparticle in the N -th shell. The subscript $\tau = \pm$ represents the two channels which are constructed by the $l = 0$ and $l = 1$ orbitals as discussed in the last part of the previous subsection. We can obtain the same NRG Hamiltonian for a $d_{x-2-y^2} + id_{xy}$ -wave. In order to check reliability of numerical results, we compare different data

by changing cutoff of the eigenstates and the discretization parameter Λ . In the NRG results we show throughout this paper, we keep the lowest-lying ~ 500 states at each renormalization step and take $\Lambda = 3$. For calculating a local spin susceptibility, we keep ~ 1200 states to reduce the cutoff dependence.

III. KONDO EFFECT IN P -WAVE SUPERCONDUCTOR

In this section we present our results for the Kondo effect in the $p_x + ip_y$ -wave superconducting state. We show numerical results and then discuss the strong coupling limit to see how the orbital dynamics of the Cooper pairs affects the Kondo singlet.

A. Numerical results

First we discuss a $\tilde{\Delta} = 0$ case where the only $l = 0$ electrons couple with the local spin and the $l = 1$ electrons are free from the impurity. The NRG result shows that the Kondo singlet is realized by the relevant $l = 0$ electrons at the stable fixed point. This fixed point is of strong coupling type ($|\tilde{J}| \rightarrow \infty$) and the fixed-point NRG energy levels is reproduced by truncating the $N = 0$ site (impurity site) from the $l = 0$ chain of the conduction electrons.²⁹ On the other hand, the fixed point for $l = 1$ is of free-electron type ($\tilde{J} = 0$). One of the two chains produces a zero energy state in one-particle energy levels at the fixed point. We can fill at most two particles (spin up and down) into the zero energy state and we have quartet ground state for $\tilde{\Delta} = 0$.

Next we discuss a $\tilde{J} = \tilde{V} = 0$ case with finite $\tilde{\Delta}$. In this case, the NRG Hamiltonian (2.23) consists of the two independent $\tau = \pm$ channels related to the superconducting state. The Hamiltonian of each channel is same as that of the free s -wave superconductivity.⁸ At the odd renormalization steps, one-particle energy level for the free s -wave is particle-hole symmetric. At the even steps, there is an additional one-particle energy level which is hole-like (particle-like) for a positive (negative) $\tilde{\Delta}$. In the present $p_x + ip_y$ -wave case, one particle energy level is symmetric at all the renormalization steps, since the signs of $\tilde{\Delta}$ for the $\tau = \pm$ channels are different each other. In one-particle energy levels, there is no zero energy state due to the finite $\tilde{\Delta}$. The ground state is obtained by filling particles into the energy levels and we have a single ground state.

Let us turn to the Kondo problem in the $p_x + ip_y$ -wave state. In this case both $\tilde{\Delta}$ and $\tilde{J}$ are finite. Since $\tilde{J}$ grows up as $\Lambda^{N/2}$ with increasing N , it is relevant in the Kondo effect. In addition, $\tilde{\Delta}$ is also relevant since it grows as in Eq. (2.23). Therefore we can expect a competition between the Kondo effect and the superconductivity.⁸ As




FIG. 1. T_K/Δ dependence of the bound state energy $(E_B/\Delta)^*$ for $p_x + ip_y$ -wave. The ground state is always a spin doublet and the bound state level is measured from the ground state for each T_K/Δ . The first excited state is a particle-hole doublet with no spin for $\tilde{V} = 0$. The circle and star represent the result with fixed T_K for $T_K = 9.20 \times 10^{-5}$ and 1.76×10^{-3} , respectively. The triangle-up and triangle-down are the results for fixed $\tilde{\Delta} = 0.001$ and 0.005 , respectively. Here $\tilde{\Delta} = 1.5\Delta$. For both T_K and Δ , a unit of the energy is the band width $2E_F$.

the renormalization step increases, the excited state energies increase with $\tilde{\Delta}_N \sim \Lambda^{N/2}$, which is the energy gap at the N -th renormalization step. Since the magnetic impurity can destroy the superconductivity, we have bound states below $\tilde{\Delta}_N$. The ratio of the bound state energy and $\tilde{\Delta}_N$ approaches a constant value with the increase of the renormalization step. We express the convergent value of the ratio by $(E_B/\Delta)^*$. Figure 1 shows the T_K/Δ dependence of the first excited state (bound state) for $\tilde{V} = 0$, where the Kondo temperature is defined by

$$T_K = \sqrt{|\tilde{J}|} \exp(-1/|\tilde{J}|). \quad (3.1)$$

Here, a unit of the energy is taken to be the band width $2E_F$. We find that the bound state energy level is scaled by T_K/Δ as in the s -wave case. In a small T_K/Δ region, the ground state is a spin doublet as expected, and the first excited state is a particle-hole doublet with no spin. With the increase of T_K/Δ , the energy level of the bound state (the first excited state) decreases (see Fig. 1). As $T_K/\Delta \rightarrow \infty$, it approaches the ground state energy and the ground state is four-fold degenerate. The result for $T_K/\Delta \rightarrow \infty$ is connected smoothly to that for $\Delta = 0$.

The most important point in Fig. 1 is that the ground state is always a spin doublet. This means that the spin of the ground state survives even in the strong coupling limit for the $p_x + ip_y$ -wave superconductivity. In the s -wave case, the Kondo singlet is realized in the large T_K/Δ region⁸, resulting in a spin singlet ground state. Thus the Kondo effect for the $p_x + ip_y$ -wave is quite different from that for the conventional s -wave.

While the ground state is always a spin doublet for the $p_x + ip_y$ -wave, we find that the T_K/Δ dependence of the bound state energy level is very similar to that of the s -wave. For the s -wave, the Kondo singlet is real-

ized in $T_K/\Delta > 0.3$ region against the superconducting energy gap.⁸ In Fig. 1, the curvature of the $(E_B/\Delta)^*$ curve found around $(E_B/\Delta)^* = 0.5$ is very similar to that of the s -wave case found with the interchange of the ground state. This represents a crossover from the Δ dominant region to the T_K dominant region. It occurs around $T_K/\Delta = 0.3$ as in the s -wave case. Thus the Kondo effect can overcome the energy gap also in the $p_x + ip_y$ -wave case, although the ground state is still a spin doublet. This means that the competition between the Kondo effect and the energy gap is characterized by such T_K/Δ dependence. All the above results hold for the $d_{x^2-y^2} + id_{xy}$ -wave as well, since we can obtain the same NRG Hamiltonian.

Finally, we mention the effect of the potential scattering $\tilde{V}$ in Eq. (2.23). It is well known that a single non-magnetic impurity destroys the unconventional superconductivity. It can generate a bound state locally around the impurity. If we treat the magnetic impurity realistically, we have to start from the Anderson model. Since the generalized Kondo model consists of both magnetic and non-magnetic interactions, we have to take both couplings into account. Since the effect of $\tilde{V}$ breaks the particle-hole symmetry, it lifts the degeneracy of the first excited particle-hole doublet. However, the ground state is still a spin doublet and $\tilde{V}$ is irrelevant even if it is large enough.

B. Strong coupling limit

Here we study the mechanism of the spin doublet ground state for the $p_x + ip_y$ -wave. Let us discuss the strong coupling limit ($|\tilde{J}| \rightarrow \infty$) case. The Hamiltonian H_0 in Eq. (2.23) can exhibit the solution in this limit. Returning the channel (τ) representation to the angular momentum (l) one, we can rewrite the H_0 for $\tilde{V} = 0$ as

$$\begin{aligned} H_0 &= H_{0J} + H_{0\Delta}, \\ H_{0J} &= -\tilde{J} \sum_{\sigma\sigma'} \mathbf{S} \cdot \boldsymbol{\sigma}_{\sigma\sigma'} g_{0\sigma}^\dagger g_{0\sigma'}, \\ H_{0\Delta} &= -\tilde{\Delta} \sum_{\sigma} (g_{0\sigma}^\dagger g_{1\sigma} + g_{1\sigma}^\dagger g_{0\sigma}), \end{aligned} \quad (3.2)$$

where the following transformation has been used:

$$c_{N=0,\tau,\sigma} = \frac{1}{\sqrt{2}} (g_{l=0,\sigma} + \tau g_{l=1,\sigma}). \quad (3.3)$$

Here $g_{l=0,\sigma}$ and $g_{l=1,\sigma}$ correspond to $f_{00\sigma}$ and $f_{01\sigma}^\dagger$, respectively. Notice that the order parameter corresponds to the hopping parameter between the $l = 0$ and $l = 1$ local orbital sites, since the $p_x + ip_y$ -wave Cooper pair consists of the $l = 0$ and $l = 1$ particles. When $|\tilde{J}| \rightarrow \infty$, the $l = 0$ particle couples with the local spin strongly to form a spin singlet $|s\rangle$:

$$|s\rangle = \frac{1}{\sqrt{2}} (g_{0\uparrow}^\dagger |\downarrow\rangle - g_{0\downarrow}^\dagger |\uparrow\rangle), \quad (3.4)$$




FIG. 2. Excited states generated from the unperturbed ground ones by the perturbation of $\tilde{\Delta}$. (a) Particle-hole doublet case. (b) Spin doublet case.

where $|\uparrow\rangle$ and $|\downarrow\rangle$ represent the local spin states. Since the hopping is forbidden, the ground states are 4-fold degenerate for $|J| \rightarrow \infty$:

$$\begin{aligned} |s \uparrow\rangle &= g_{1\uparrow}^\dagger |s\rangle, \\ |s \downarrow\rangle &= g_{1\downarrow}^\dagger |s\rangle, \\ |p1\rangle &= |s\rangle, \\ |p2\rangle &= g_{1\uparrow}^\dagger g_{1\downarrow}^\dagger |s\rangle, \end{aligned} \quad (3.5)$$

where $|s\sigma\rangle$ ($\sigma = \uparrow, \downarrow$) and $|pi\rangle$ ($i = 1, 2$) are the spin doublet and particle-hole doublet, respectively. At a finite $|\tilde{J}|$, $\tilde{\Delta}$ in Eq. (3.2) lifts the degeneracy. The perturbation (the $\tilde{\Delta}$ term) generates excited states from the ground states in Eq. (3.5). We show the connection among these states in Fig. 2. One of the particle-hole doublet states is connected to a single excited state. For $|p1\rangle$, the excited state is given by [see Fig. 2 (a)]:

$$|p1'\rangle = \frac{1}{\sqrt{2}} (g_{1\uparrow}^\dagger |\downarrow\rangle - g_{1\downarrow}^\dagger |\uparrow\rangle). \quad (3.6)$$

The similar argument holds for $|p2\rangle$. We can obtain the following eigenvalue for the particle-hole doublet up to the fourth order of the perturbation:

$$E_{S=0} = \left[-\frac{3}{2} - \frac{2}{3} \left(\frac{\tilde{\Delta}}{|\tilde{J}|} \right)^2 + \frac{8}{27} \left(\frac{\tilde{\Delta}}{|\tilde{J}|} \right)^4 \right] |\tilde{J}|. \quad (3.7)$$

For one of the spin doublet states, it is connected to four excited states by the perturbation up to the fourth order. For $|s \uparrow\rangle$, these excited states are given as follows [see Fig. 2 (b)]:

$$\begin{aligned} |0 \uparrow\rangle &= g_{0\uparrow}^\dagger g_{0\downarrow}^\dagger |\uparrow\rangle, \\ |1 \uparrow\rangle &= g_{1\uparrow}^\dagger g_{1\downarrow}^\dagger |\uparrow\rangle, \\ |t1\rangle &= g_{1\downarrow}^\dagger g_{0\uparrow}^\dagger |\uparrow\rangle, \\ |t2\rangle &= \frac{1}{\sqrt{2}} g_{1\uparrow}^\dagger [g_{0\uparrow}^\dagger |\downarrow\rangle + g_{0\downarrow}^\dagger |\uparrow\rangle], \end{aligned} \quad (3.8)$$

and the similar connection is obtained for $|s \downarrow\rangle$ as well. The energy eigenvalue can be obtained from the following equation:




FIG. 3. Schematic picture of the ground state wavefunction. (a) Weak coupling limit ($T_K/\Delta \rightarrow 0$). (b) Strong coupling limit ($T_K/\Delta \rightarrow \infty$).

$$\begin{aligned} e(e-1/2) [e^3 + e^2 - (4x^2 + 3/4)e - 4x^2] &= 0, \\ e = E/|\tilde{J}|, \quad x = |\tilde{\Delta}|/|\tilde{J}|. \end{aligned} \quad (3.9)$$

After expanding the lowest e in x up to the fourth order, we obtain the energy for the spin doublet as

$$E_{S=1/2} = \left[-\frac{3}{2} - \frac{2}{3} \left(\frac{\tilde{\Delta}}{|\tilde{J}|} \right)^2 - \frac{10}{27} \left(\frac{\tilde{\Delta}}{|\tilde{J}|} \right)^4 \right] |\tilde{J}|. \quad (3.10)$$

The energy difference between the two doublets is given by

$$E_{S=0} - E_{S=1/2} = \frac{2}{3} |\tilde{J}| \left(\frac{\tilde{\Delta}}{|\tilde{J}|} \right)^4. \quad (3.11)$$

Thus, the order parameter $\tilde{\Delta}$ stabilizes the spin doublet even if it is very small.

In the s -wave case, the superconducting part of H_0 is given by

$$H_{0\Delta} = -\tilde{\Delta} (g_{0\uparrow}^\dagger g_{0\uparrow} + g_{0\downarrow}^\dagger g_{0\downarrow} - 1). \quad (3.12)$$

We notice that it is not given by hopping between the sites but is given by a potential on the $l=0$ site. This is because the s -wave Cooper pair is formed by only the $l=0$ quasiparticles. In the $|\tilde{J}| \rightarrow \infty$ limit, we have a Kondo singlet ground state as expected.

Comparing the $p_x + ip_y$ -wave result with the s -wave one, we conclude that the spin of the ground state is realized by the orbital dynamics of the $p_x + ip_y$ -wave Cooper pairs. This point is illustrated by Fig. 3. In a weak coupling region, the spin of the ground state is mainly produced by the local spin itself [see Fig. 3(a)]. In the strong coupling limit, however, a Kondo singlet is formed by the $l=0$ conduction electrons. Therefore the local spin is almost quenched by the $l=0$ electrons. In this limit, one of the paired electrons having $l=1$




FIG. 4. (a) Temperature dependence of $T\chi_{\text{local}}$ for various $\tilde{\Delta}$. Here $\tilde{\Delta} = 1.5\Delta$. The exchange coupling is fixed as $\tilde{J} = -0.25$ ($T_K = 5.34 \times 10^{-4}$). The unit is $(g_0\mu_B)^2$, where g_0 is a bare g -factor of the local spin. (b) T_K/Δ dependence of the effective g -factor. The circle, star, and triangle-up represent the results with fixed $T_K = 9.20 \times 10^{-5}$, 5.34×10^{-4} , and 1.76×10^{-3} , respectively. For both T_K and Δ , a unit of the energy is the band width $2E_F$.

angular momentum is connected weakly with the Kondo singlet to gain the superconducting condensation energy. The latter electrons generate the ground state spin as shown in Fig. 3(b).

C. Impurity susceptibility

Next we see how the spin of the magnetic impurity is screened by the Kondo effect. For this purpose we calculate a local spin susceptibility which is defined by

$$\chi_{\text{local}}(T) = g\mu_B \frac{\langle S_z \rangle}{h} \Big|_{h \rightarrow 0}, \quad (3.13a)$$

$$\langle S_z \rangle = \frac{\text{Tr} [S_z \exp(-\bar{\beta} H_N)]}{\text{Tr} [\exp(-\bar{\beta} H_N)]}. \quad (3.13b)$$

Here S_z is the local spin operator, and $\bar{\beta} \sim 1$ is taken in the calculation. The temperature T is given by²⁹⁻³¹

$$T = \frac{1}{2}(1 + \Lambda^{-1})\Lambda^{-(N-1)/2}/\bar{\beta}. \quad (3.14)$$

A small magnetic field h is applied only at the local spin site. The impurity part of the Hamiltonian H_0 in Eq. (2.23b) is replaced as




FIG. 5. Bound state energy calculated by the NRG method for the Ising case. It is normalized by Δ . The solid line is the analytic solution given by $E_B/\Delta = [1 + (\alpha\tilde{J}_z)^2]^{-1/2}$, where $\tilde{J}_z = 1.5\tilde{J}_z$, and α is a constant. We have used $\alpha = 0.465$ to fit the NRG result.

$$H_0 \rightarrow H_0 - g\mu_B S_z h \Lambda^{-1/2}. \quad (3.15)$$

Therefore χ_{local} represents magnetic response of the local spin. In Fig. 4(a) we show the temperature dependence of χ_{local} for various Δ with a fixed exchange coupling $\tilde{J}$. We can see that $T\chi_{\text{local}}$ decreases monotonically as we decrease temperature (increase the renormalization step) for each Δ , and that it becomes constant at low temperatures, indicating the Curie law. However, the effective g -factor of the local spin depends on both $\tilde{J}$ and Δ shown in Fig. 4(b). We can see that g is scaled by T_K/Δ as in the case of the energy of the first excited state [see Fig. 1]. This result indicates that the local spin is actually screened by the Kondo effect, although the ground state is a spin doublet.

In order to see the quantum effect (Kondo effect) of the local spin, we compare the result of the bound state energy for a Ising (classical) spin [see Fig. 5]. The bound state energy position is given by

$$E_B = \frac{\Delta}{\sqrt{1 + \gamma^2}}, \quad (3.16a)$$

$$\gamma = \frac{\pi N_0}{4\Omega} |J_z|, \quad (3.16b)$$

which is essentially same as in the non-magnetic impurity case.²⁵ Here N_0 is the density of state at the Fermi energy in the normal state. Ω is the volume of the system. The bound state energy decreases as we increase the Ising coupling $|J_z|$. This point is similar to the result of the quantum spin case shown in Fig. 1. However, the bound state energy is independent of Δ for the Ising spin, while it is a function of T_K/Δ for the quantum spin. In addition, we have to use an extremely larger $|J_z|$ value ($|J_z| \gg 1$) for the Ising spin to obtain a small $(E_B/\Delta)^*$. This indicates that the spin-flip effect (quantum effect) is essential to our result, and that the Kondo effect actually occurs in the $p_x + ip_y$ -wave superconductor.

IV. CONCLUSION AND DISCUSSION

We have studied the novel Kondo effect due to the orbital effect of the Cooper pairs of unconventional superconductors. In particular, we have focused on the $p_x + ip_y$ -wave ($d_{x^2-y^2} + id_{xy}$ -wave) pairing state. Regarding the magnetic impurity, we have assumed an s -wave (short-range type) impurity scattering, which is the simplest but is primarily important in real materials. The characteristic point of the unconventional superconductor is that the Cooper pair has orbital degrees of freedom. Therefore the superconducting quasiparticles consist of electrons with various angular momenta. The object of this paper is to clarify how the orbital dynamics of the Cooper pairs affects the Kondo effect in unconventional superconductors. The followings are our main results.

(1) $p_x + ip_y$ -wave

In the superconducting state, the low-energy excitations are moved out of the superconducting energy gap. The $p_x + ip_y$ -wave superconductor is a fully gapped system. Unlike the s -wave superconductor, the $p_x + ip_y$ -wave Cooper pairs have the orbital degrees of freedom. The local spin couples with only $l = 0$ electrons which are paired with the $l = 1$ ones. However, the $l = 1$ electrons can interact with the local spin via the superconducting order parameter. We note that there is no direct coupling between the local spin and the $l = 1$ electrons. In contrast to the two-channel Kondo effect derived from the atomic structure of the impurity,¹ the present two-channel property is the consequence of the internal degrees of freedom of the conduction electron system. Our NRG result shows that the ground state is a spin doublet over all the T_K/Δ region. This is completely different from the result of the s -wave superconductor in which the Kondo singlet is realized in a large T_K/Δ region. We also find that the first excited energies are scaled by T_K/Δ . In the strong coupling limit ($T_K/\Delta \rightarrow \infty$), it is found that the effective g -factor of the local spin is strongly reduced by the Kondo effect, while the $p_x + ip_y$ -wave Cooper pair couples with the Kondo singlet, generating the spin of the ground state. Thus the orbital effect of the Cooper pair is important in the Kondo effect we have discussed here.

(2) Origins of the multichannel property

Generally, there are two origins of the multichannel property in the Kondo effect. The first one is the internal degrees of freedom of the impurity atom as pointed out by Nozières and Blandin.¹ In this case, the channels of the conduction electron system is controlled by the atomic structure of the impurity atom.

The second one is related to the conduction electron systems. In this paper we have investigated the Kondo effect in the unconventional superconductors. As shown in the $p_x + ip_y$ -wave case, the multichannel property comes from the angular momenta of the Cooper pairs. The independent channels of the superconducting quasiparticles couple with each other at the impurity even if there is no

internal degrees of freedom in the impurity atom (just a local spin). This is because the symmetry of the conduction electron system is different from that of the impurity scattering. As a different variety besides the superconducting systems, we can introduce strong momentum dependence into the bare conduction band. For example, a singular pseudo-gap in the vicinity of the Fermi energy produces unusual Kondo behavior.³⁵ Thus the internal degrees of freedom of both the impurity and the conduction electron system are important for the Kondo effect in real materials.

ACKNOWLEDGMENTS

We are grateful to S. Curnoe, H. Kusunose, O. Sakai, H. Shiba, R. Shiina, and K. Ueda for helpful discussions and comments.

APPENDIX A: ANOTHER DERIVATION OF THE NRG HAMILTONIAN

In Appendix, we present another derivation of the NRG Hamiltonian (2.23), following Satori *et al.*⁸ First, we apply the logarithmic discretization to the conduction band. Then we obtain the hopping type of Hamiltonian,

$$\bar{H}_{\text{kin}} = \frac{1 + \Lambda^{-1}}{2} \times \sum_{n=0}^{\infty} \sum_{l\sigma} \Lambda^{-n/2} \varepsilon_n (f_{nl\sigma}^\dagger f_{n+1,l\sigma} + f_{n+1,l\sigma}^\dagger f_{nl\sigma}), \quad (\text{A1})$$

for the free electrons. Here $f_{nl\sigma}^\dagger$ and $f_{nl\sigma}$ are fermion operators corresponding to the conduction electrons with angular momentum l . For the BCS pairing interaction in the $p_x + ip_y$ -wave superconductors, we can also obtain the following Hamiltonian:

$$\bar{H}_\Delta = \sum_{n=0}^{\infty} \sum_{\sigma} (-i\bar{\Delta}) (f_{n,0,\sigma}^\dagger f_{n,1,-\sigma}^\dagger - f_{n,1,-\sigma} f_{n,0,\sigma}). \quad (\text{A2})$$

To diagonalize $\bar{H}_\Delta$, we use the following Bogoliubov transformation:

$$b_{n,+,\sigma}^\dagger = \frac{1}{\sqrt{2}} (f_{n,0,\sigma}^\dagger - i f_{n,1,-\sigma}), \quad (\text{A3a})$$

$$b_{n,-,-\sigma} = \frac{1}{\sqrt{2}} (f_{n,1,-\sigma} - i f_{n,0,\sigma}^\dagger), \quad (\text{A3b})$$

and we obtain

$$\bar{H}_\Delta = -\bar{\Delta} \sum_{n=0}^{\infty} \left(\sum_{\tau\sigma} b_{n\tau\sigma}^\dagger b_{n\tau\sigma} - 2 \right). \quad (\text{A4})$$

The impurity Hamiltonian is expressed by the Bogoliubov operators $b_{n\tau\sigma}^\dagger$ and $b_{n\tau\sigma}$ as

$$\begin{aligned}\bar{H}_{\text{imp}} &= -\bar{J} \sum_{\sigma\sigma'} \mathbf{S} \cdot \boldsymbol{\sigma}_{\sigma\sigma'} f_{00\sigma}^\dagger f_{00\sigma'} \\ &= -\frac{\bar{J}}{2} \sum_{\sigma\sigma'} \mathbf{S} \cdot \boldsymbol{\sigma}_{\sigma\sigma'} (b_{0,+,\sigma}^\dagger + ib_{0,-,-\sigma}) \\ &\quad \times (b_{0,+,\sigma'} - ib_{0,-,-\sigma'}). \quad (\text{A5})\end{aligned}$$

The Hamiltonian for the conduction electrons is given by

$$\begin{aligned}\bar{H}_{\text{kin}} &= \frac{1 + \Lambda^{-1}}{2} \sum_{n=0}^{\infty} \sum_{\sigma} \Lambda^{-n/2} \varepsilon_n(-i) \\ &\quad \times (b_{n,+,\sigma} b_{n+1,-,-\sigma} + b_{n,+,\sigma}^\dagger b_{n+1,-,-\sigma}^\dagger \\ &\quad + b_{n+1,+,\sigma} b_{n,-,-\sigma} + b_{n+1,+,\sigma}^\dagger b_{n,-,-\sigma}^\dagger). \quad (\text{A6})\end{aligned}$$

Next we apply the following particle-hole transformation:

$$\begin{aligned}c_{2n,+,\sigma}^\dagger &= b_{2n,+,\sigma}^\dagger, \quad c_{2n,-,\sigma}^\dagger = ib_{2n,-,-\sigma}, \\ c_{2n-1,+,\sigma}^\dagger &= -ib_{2n-1,-,-\sigma}, \quad c_{2n-1,-,\sigma}^\dagger = -b_{2n-1,+,\sigma}^\dagger. \quad (\text{A7})\end{aligned}$$

In this manner we can exchange the roles of particles and holes to keep the number of fermion quasiparticles. Then we obtain the same form of the Hamiltonian given in (2.23). The above argument can be applied to the case of $d_{x^2-y^2} + id_{xy}$ -wave superconductors by replacing $i\Delta$ to $-\sigma\Delta$. In this case, the Bogoliubov transformation is given by

$$b_{n,+,\sigma}^\dagger = \frac{1}{\sqrt{2}} (f_{n,0,\sigma}^\dagger + \sigma f_{n,2,-\sigma}), \quad (\text{A8a})$$

$$b_{n,-,-\sigma} = \frac{1}{\sqrt{2}} (f_{n,2,-\sigma} - \sigma f_{n,0,\sigma}^\dagger), \quad (\text{A8b})$$

and the roles of particles and holes are exchanged by the following transformation:

$$\begin{aligned}c_{2n,+,\sigma}^\dagger &= b_{2n,+,\sigma}^\dagger, \quad c_{2n,-,\sigma}^\dagger = -\sigma b_{2n,-,-\sigma}, \\ c_{2n-1,+,\sigma}^\dagger &= -\sigma b_{2n-1,-,-\sigma}, \quad c_{2n-1,-,\sigma}^\dagger = b_{2n-1,+,\sigma}^\dagger. \quad (\text{A9})\end{aligned}$$

-
- ¹ P. Nozières and A. Blandin, *J. Physique* **41**, 193 (1980).
² For a review, D. L. Cox and A. Zawadowski, *Adv. Phys.* **47**, 599 (1998).
³ H. Shiba, *Prog. Theor. Phys.* **40**, 435 (1968).
⁴ A. Sakurai, *Prog. Theor. Phys.* **44**, 1472 (1970).
⁵ E. Müller-Hartmann, *Magnetism*, ed. H. Shul, Vol. 5, p. 353 (Academic, 1973); E. Müller-Hartmann and J. Zittartz, *Z. Phys.* **234**, 58 (1970).
⁶ T. Matsura, *Prog. Theor. Phys.* **57**, 1823 (1977).
⁷ M. Jarrell, D. S. Sivia, and B. Patton, *Phys. Rev. B* **42**, 4804 (1990).
⁸ K. Satori, H. Shiba, O. Sakai, and Y. Shimizu, *J. Phys. Soc. Jpn.* **61**, 3239 (1992).
⁹ O. Sakai, Y. Shimizu, H. Shiba, and K. Satori, *J. Phys. Soc. Jpn.* **62**, 3181 (1993).
¹⁰ T. Yoshioka and Y. Ohashi, *J. Phys. Soc. Jpn.* **67**, 1332 (1998); *ibid.* **69**, 1812 (2000).
¹¹ S. Schmitt-Rink, K. Miyake, and C. M. Varma, *Phys. Rev. Lett.* **57**, 2575 (1986); P. J. Hirschfeld and N. Goldenfeld, *Phys. Rev. B* **48**, 4219 (1993).

- ¹² K. Ishida, Y. Kitaoka, N. Ogata, T. Kamino, K. Asayama, J. R. Cooper, *J. Phys. Soc. Jpn.* **62**, 2803 (1993).
¹³ S. H. Pan, E. W. Hudson, K. M. Lang, H. Eisaki, S. Uchida, and J. C. Davis, *Nature (London)* **403**, 746 (2000).
¹⁴ C.-R. Hu, *Phys. Rev. Lett.* **72**, 1526 (1994).
¹⁵ I. Iguchi, W. Wang, M. Yamazaki, Y. Tanaka, and S. Kashiwaya, *Phys. Rev. B* **62**, R6131 (2000).
¹⁶ H. Alloul, P. Mendels, H. Casalta, J. F. Marucco, and J. Arabski, *Phys. Rev. Lett.* **67**, 3140 (1991); A. V. Mahajan, H. Alloul, G. Collin, and J. F. Marucco, *ibid.* **72**, 3100 (1994); Y. Sidis, P. Bourges, H. F. Fong, B. Keimer, L. P. Regnault, J. Bossy, A. Ivanov, B. Hennion, P. Gautier-Picard, G. Collin, D. L. Millius, and I. A. Aksay, *ibid.* **84**, 5900 (2000).
¹⁷ D. L. Sisson, S. G. Doettinger, A. Kapitulnik, R. Liang, D. A. Bonn, and W. N. Hardy, *Phys. Rev. B* **61**, 3604 (2000).
¹⁸ J. Bobroff, H. Alloul, W. A. MacFarlane, P. Mendels, N. Blanchard, G. Collin, and J.-F. Marucco, *Phys. Rev. Lett.* **86**, 4116 (2001).
¹⁹ M. E. Simon and C. M. Varma, *Phys. Rev. B* **60**, 9744 (1999).
²⁰ Y. Maeno, H. Hashimoto, K. Yoshida, S. Nishizaki, T. Fujita, J. G. Bednorz, and F. Lichtenberg, *Nature (London)* **372**, 532 (1994).
²¹ K. Ishida, H. Mukuda, Y. Kitaoka, K. Asayama, Z. Q. Mao, Y. Mori, and Y. Maeno, *Nature (London)* **396**, 658 (1998).
²² T. M. Rice and M. Sigrist, *J. Phys. Condens. Matter*, **7**, L643 (1995); G. Baskaran, *Physica (Amsterdam)* **223B-224B**, 490 (1996); M. Sigrist, D. F. Agterberg, A. Furusaki, C. Honerkamp, K. K. Ng, T. M. Rice, and M. E. Zhitomirsky, *Physica (Amsterdam)* **317C-318C**, 134 (1999).
²³ G. M. Luke, Y. Fudamoto, K. M. Kojima, M. I. Larkin, J. Merrin, B. Nachumi, Y. J. Uemura, Y. Maeno, Z. Q. Mao, Y. Mori, H. Nakamura, and M. Sigrist, *Nature (London)* **394**, 558 (1998).
²⁴ M. Matsumoto and M. Sigrist, *J. Phys. Soc. Jpn.* **68**, 994 (1999); *ibid.* **68**, 3120 (1999).
²⁵ Y. Okuno, M. Matsumoto, and M. Sigrist, *J. Phys. Soc. Jpn.* **68**, 3054 (1999).
²⁶ M. Matsumoto and M. Sigrist, *Physica (Amsterdam)* **284B-288B**, 545 (2000).
²⁷ J. Goryo and K. Ishikawa, *Phys. Lett. A* **260**, 294 (1999); A. Furusaki, M. Matsumoto, and M. Sigrist, *Phys. Rev. B* **64**, 054514 (2001).
²⁸ J. Goryo, *Phys. Rev. B* **61**, 4222 (2000); M. Matsumoto and M. Sigrist, *Physica (Amsterdam)* **281B-282B**, 973 (2000); M. Matsumoto and R. Heeb, *cond-mat/0101155*.
²⁹ K. G. Wilson, *Rev. Mod. Phys.* **47**, 773 (1975).
³⁰ H. R. Krishna-Murthy, J. W. Wilkins, and K. G. Wilson, *Phys. Rev. B* **21**, 1003 (1980); *ibid.* **21**, 1044 (1980).
³¹ O. Sakai, Y. Shimizu, and T. Kasuya, *Prog. Theor. Phys. Suppl.* **108**, 73 (1992).
³² D. M. Cragg, P. Loyd, and P. Nozières, *J. Phys. C* **13**, 803 (1980).
³³ H. B. Pang and D. L. Cox, *Phys. Rev. B* **44**, 9454 (1991).
³⁴ D. L. Cox, *Phys. Rev. Lett.* **59**, 1240 (1987).
³⁵ D. Withoff and E. Fradkin, *Phys. Rev. Lett.* **64**, 1835 (1990); C. Gonzalez-Buxton and K. Ingersent, *Phys. Rev. B* **57**, 14254 (1998).